

Ward et al. (2010) recommended that “the relationship between occupational exposure to carbon black and validated biomarkers of oxidative stress should be examined.” Despite the appeal of biomarkers of oxidative stress in pinpointing inflammatory changes associated with malignant and nonmalignant illnesses, such markers are nonspecific, not well validated, and appear not “ready for prime time,” as noted in a recent symposium on nanotoxicology (Fischman et al. 2011).

A meta-analysis of all three major carbon black cohorts (United States, United Kingdom, and Germany) to assess risk of heart disease is also under way. In a recent position paper, Brook et al. (2004) noted that particle exposure may play a role in the development of heart disease.

Ward et al. (2010) suggested evaluating carbon black particle size and surface area. However, the physical and chemical properties of untreated manufactured carbon blacks are distinctly different from ubiquitous carbon core particulates in both occupational and ambient atmospheres (Kuhlbusch and Fissan 2006). Approximately 90% of manufactured carbon black is used for tire and automotive rubber products. In products, such as toners, plastics, and surface coatings, carbon black is matrix-bound, and not an exposure risk to end-users. Care should be taken when applying quantitative models that claim to address the particle size and surface area topics (Tomenson and Morfeld 2010).

The authors serve as scientific advisors to the International Carbon Black Association (ICBA), a scientific, non-profit corporation originally founded in 1977, with the purpose of sponsoring, conducting, and participating in investigations, research, and analyses relating to the health, safety, and environmental aspects of the production and use of carbon black. This manuscript was neither influenced by the ICBA nor by any company funding the ICBA, nor does it present any view or opinion of the ICBA or of the companies. H.M. is president of Muranko and Associates, a consulting company that provides industrial hygiene and safety services and serves as expert witness in industrial safety cases.

Robert J. McCunney

Massachusetts Institute of Technology
Cambridge, MA
E-mail: mccunney@MIT.EDU

Peter Morfeld

Institute for Occupational Medicine
at Cologne University
Cologne, Germany

Len Levy

Emeritus Professor
Institute of Environment and Health
Cranfield University
Cranfield, Bedfordshire, United Kingdom

Henry Muranko

Muranko and Associates
Scottsdale, Arizona

The Scientific Advisory Group of the International Carbon Black Association (ICBA) gave helpful comments on an earlier version of this letter.

REFERENCES

- Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, et al. 2004. Air pollution and cardiovascular disease: a statement for healthcare professionals from the Expert Panel on Population and Prevention Science of the American Heart Association. *Circulation* 109(21):2655–2671.
- Dell LD, Mundt KA, Luippold RS, Nunes AP, Cohen L, Burch MT, et al. 2006. A cohort mortality study of employees in the U.S. carbon black industry. *J Occup Environ Med* 48(12):1219–1229.
- Fischman M, Storey E, McCunney RJ, Kosnett M. 2011. National Institute for Occupational Safety and Health Nanomaterials and Worker Health Conference—medical surveillance session summary report. *J Occup Med* 53(6 suppl):S35–S37.
- Kuhlbusch TA, Fissan H. 2006. Particle characteristics in the reactor and pelletizing areas of carbon black production. *J Occup Environ Hyg* 3(10):558–567.
- Morfeld P, McCunney RJ. 2007. Carbon black and lung cancer: testing a new exposure metric in a German cohort. *Am J Ind Med* 50(8):565–567.
- Morfeld P, McCunney RJ. 2009. Carbon black and lung cancer—testing a novel exposure metric by multi-model inference. *Am J Ind Med* 52(11):890–899.
- Morfeld P, McCunney R. 2010. Bayesian bias adjustments of the lung cancer SMR in a cohort of German carbon black production workers. *J Occup Med Toxicol* 5:23; doi:10.1186/1745-6673-5-23 [11 August 2010].
- Sorahan T, Harrington JM. 2007. A “lugged” analysis of lung cancer risks in UK carbon black production workers, 1951–2004. *Am J Ind Med* 50(8):555–564.
- Tomenson JA, Morfeld P. 2010. Multiple errors made by authors result in a huge overestimation of potential exposure to particles in the size range 10–30 nm in TiO₂ nanoparticle production facilities [Letter]. *J Hazard Mater* 183:954–955.
- Ward EM, Schulte PA, Straif K, Hopf NB, Caldwell JC, Carréon T, et al. 2010. Research recommendations for selected IARC-classified agents. *Environ Health Perspect* 118:1355–1362.

Carbon Black: Kuempel et al. Respond

doi:10.1289/ehp.1103444R

We appreciate the comments and additional information from McCunney et al. We are pleased to learn of the new epidemiological studies that are under way in the U.S. and E.U. carbon black cohorts. These studies may provide the opportunity to fill some of the research gaps discussed in our review (Ward et al. 2010). As mentioned by McCunney et al. in their letter, we recommend the collection of particle size-specific and quantitative exposure data, and the recruitment of additional facilities (Ward et al.). Studies in animals have shown relationships between the particle surface area dose of poorly soluble particles (including carbon black) in the lungs and biomarkers of oxidative stress and inflammation in rats and mice (Elder et al. 2005; Sager and Castranova 2009; Stoeger et al. 2006) and lung tumors in rats (Driscoll 1996; Heinrich et al. 1995; Nikula et al. 1995). Although these relationships

with particle surface area dose have not been reported in human studies, exposure to carbon black by particle mass has been associated with respiratory effects including lung function decrements in workers (Gardiner et al. 2001).

Concerning biomarkers of oxidative stress, we think the epidemiology studies under way may provide an opportunity to investigate and test hypotheses about possible biomarkers of exposure and response to carbon black. As we discussed in our paper (Ward et al. 2010), although oxidative stress has been invoked as a mechanism in the carcinogenicity of a number of agents (including particles such as carbon black), methodological challenges to the validation of oxidative stress biomarker assays remain. To facilitate this process, guidelines have been developed to standardize the collection and measurement of oxidative stress biomarkers in humans (American Thoracic Society 1999; Horváth et al. 2005).

We look forward to further reports from the carbon black mortality studies, including exposure–response analyses, which could help fill important occupational health research gaps. Well-conducted epidemiologic studies will be particularly critical to inform carcinogen classification and risk assessment processes.

The findings and conclusions in this letter are those of the authors and do not necessarily represent the views of the National Institute for Occupational Safety and Health, the University of Birmingham, the U.S. Environmental Protection Agency, the International Agency for Research on Cancer, or the American Cancer Society Inc.

The authors declare they have no actual or potential competing financial interests.

Eileen D. Kuempel

Paul A. Schulte

National Institute for Occupational Safety and Health
Cincinnati, Ohio
E-mail: ekuempel@cdc.gov

Thomas Sorahan

Institute of Occupational and Environmental Medicine
University of Birmingham
Birmingham, United Kingdom

Jane Caldwell

U.S. Environmental Protection Agency
Research Triangle Park, North Carolina

Kurt Straif

International Agency for Research on Cancer
Lyon, France

Elizabeth Ward

American Cancer Society, Inc.
Atlanta, Georgia

REFERENCES

- American Thoracic Society. 1999. Recommendations for standardized procedures for the online and offline measurement of exhaled lower respiratory nitric oxide and nasal nitric oxide in adults and children—1999. *Am J Respir Crit Care Med* 160:2104–2117.
- Driscoll KE. 1996. Role of inflammation in the development of rat lung tumors in response to chronic particle exposure. In: *Particle Overload in the Rat Lung and Lung Cancer: Implications for Human Risk Assessment* (Mauderly JL, McCunney RJ, eds). Philadelphia:Taylor & Francis, 139–153.
- Elder A, Gelein R, Finkelstein JN, Driscoll KE, Harkema J, Oberdörster G. 2005. Effects of subchronically inhaled carbon black in three species. I. Retention kinetics, lung inflammation, and histopathology. *Toxicol Sci* 88(2):614–629.
- Gardiner K, van Tongeren M, Harrington M. 2001. Respiratory health effects from exposure to carbon black: results of the phase 2 and 3 cross sectional studies in the European carbon black manufacturing industry. *Occup Environ Med* 58(8):496–503.
- Heinrich U, Fuhst R, Rittinghausen S, Creutzenberg O, Bellmann B, Koch W, et al. 1995. Chronic inhalation exposure of Wistar rats and two different strains of mice to diesel engine exhaust, carbon black, and titanium dioxide. *Inhal Toxicol* 7:533–556.
- Horváth I, Hunt J, Barnes PF, ATS/ERS Task Force on Exhaled Breath Condensate. 2005. Exhaled breath condensate: methodological recommendations and unresolved questions. *Eur Respir J* 26:523–548.
- Nikula KJ, Snipes MB, Barr EB, Griffith WC, Henderson RF, Mauderly JL. 1995. Comparative pulmonary toxicities and carcinogenicities of chronically inhaled diesel exhaust and carbon black in F344 rats. *Fundam Appl Toxicol* 25:80–94.
- Sager TM, Castranova V. 2009. Surface area of particle administered versus mass in determining the pulmonary toxicity of ultrafine and fine carbon black: comparison to ultrafine titanium dioxide. *Part Fibre Toxicol* 6:15; doi:10.1186/1743-8977-6-15 [Online 4 May 2009].
- Stoeger T, Reinhard C, Takenaka S, Schroepel A, Karg E, Ritter B, et al. 2006. Instillation of six different ultrafine carbon particles indicates a surface area threshold dose for acute lung inflammation in mice. *Environ Health Perspect* 114:328–333.
- Ward EM, Schulte PA, Straif K, Hopf NB, Caldwell JC, Carreón T, et al. 2010. Research recommendations for selected IARC-classified agents. *Environ Health Perspect* 118:1355–1362.

for Educators

Locate lessons by topics

air
water
land
climate
food
disease

It's healthy,
it's smart,
and it's free.

www.ehponline.org/education

Stay Connected


Reading *EHP* Just Got Easier

This iPhone app provides easy access to

- » Ahead-of-Print research articles, reviews, and commentaries
- » Current issue
- » Archived issues
- » The Researcher's Perspective podcast series

You can also

- » Explore all *EHP* content using a full-text search function
- » View content as PDF or online
- » Save your favorite articles
- » E-mail articles from your iPhone



Get it for your iPhone or iPod Touch.

Get it for FREE.

Get it today.

Scan with your iPhone or iPod Touch QR reader to get the *EHP* app.



To download the free app visit the iTunes App Store, or log on to www.ehponline.org/mobile